

## Imaging in Neurological Disease of Hospitalised COVID-19 Patients



Originating in Wuhan, China, COVID-19 soon spread throughout the world and has been declared a pandemic by the World Health Organization. Italy was the second country to be hit as hard as China, with a total of 222,104 cases and 31,106 deaths (at time of publication).

A study was conducted to understand the characteristics of neurological symptoms and neuroimaging features of COVID-19 patients admitted to hospitals across Italy. This was a retrospective, multicenter study design. Three major institutions in Italy were included in the study: University of Brescia, Brescia, University of Eastern Piemonte, Novara and University of Sassari, Sassari.

### Inclusion Criteria

1. COVID-19 patients who had tested positive by rRT-PCR (real-time reverse transcriptase polymerase chain reaction) on testing of respiratory secretions obtained from endotracheal aspirates, nasopharyngeal, oropharyngeal swabs, or bronchoalveolar lavage between February 29 to April 4, 2020.
2. Acute neurological symptoms during stay at the hospital.
3. Patients who had done any neuroimaging studies such as spine or brain imaging.

### Study Results

Based on the inclusion criteria, 725 patients admitted to hospitals were reviewed, but only 108 patients actually met the criteria to be included in the study.

Total patients	108
Examined with Non-Contrast Brain CT	107 99%
Head & Neck CT angiography	17 16%
Brain MRI	20 18%

### Most common neurological symptoms

	Patients	Percentage
Altered mental status	64	59%

Ischemic stroke	34	31%
-----------------	----	-----

## Past medical history of patients

Patients with no known past medical history turned out to have the following symptoms	31/108 29%	Patients with known medical history had at least one of the following symptoms	77/108 71%
Age range (16-62)			
Acute ischaemic infarcts	10	Coronary artery disease	25 23%
Intracranial haemorrhage	2	Cerebrovascular disease	15 14%
		Hypertension	55 51%
		Diabetes	30 28%

## Findings on Brain CT

	Patients	Percentage
No acute findings	71	66%
Acute abnormalities	7	35%

The association between the age of the patient and the presence of altered mental status was statistically significant. The more aged the patient, the more prone they were to altered mental status.

## Summary of Results

1. The study showed that while there was no discernable pattern of neuroimaging characteristics, there was a predominant presence of acute ischaemic infarcts and intracranial haemorrhages.
2. Results, when compared to Wuhan, showed a lower occurrence of CNS symptoms (25% Wuhan vs. 15% Italy) but a higher occurrence of ischaemic stroke (31% Wuhan vs. 11% Italy).
3. It is not clearly understood whether the presence of neurological symptoms is due to the severity of illness or a result of a direct invasion of CNS by SARS-CoV-2. There is some evidence that ischaemic stroke in patients with severe COVID-19 might be triggered by cytokine storm syndrome.
4. There is evidence to suggest COVID-19 associated occurrence of Guillain-Barré syndrome (GBS) and some variants.

There is a strong need for neurologists and neuro-radiologists to better understand COVID-19-associated patterns in neuroimaging for better patient outcomes.

Source: [Radiology](#)

Image Credit: iStock

Published on : Sat, 13 Jun 2020