
ICU Volume 14 - Issue 2 - Summer 2014 - News

Not Enough Kidney Patients Receive Home Dialysis, Says Australian Study

Many kidney failure patients in Australia who could benefit from undergoing dialysis at home are being treated in hospitals and dialysis units, according to a study published in the Clinical Journal of the American Society of Nephrology (CJASN). This is creating significant costs for healthcare providers and causing unnecessary disruptions to patients' lives.

Home dialysis is more convenient for patients and can provide similar or better care than haemodialysis, which must be done in a clinic. Blair Grace, from the Australia and New Zealand Dialysis and Transplant Registry led a team that investigated the links between socioeconomic status and use of home dialysis (which includes peritoneal dialysis and home haemodialysis) in Australia. The researchers analysed 23,281 adult patients who started dialysis in Australia between 2000 and 2011.

Among the major findings:

- Socioeconomic status was not associated with use of home haemodialysis.
- Patients from the most advantaged areas were 37% less likely to commence peritoneal dialysis and 19% more likely to use in-centre haemodialysis than patients from the most disadvantaged areas.
- Rural areas were more disadvantaged and had higher rates of peritoneal dialysis, while privately-funded hospitals rarely used home dialysis.
- Patients from the most advantaged areas were nearly six times more likely to use private hospitals than those from the most disadvantaged areas.

"We expected to find that patients with more education and financial resources were more likely to use peritoneal dialysis and home haemodialysis, as has been demonstrated in other countries," said Dr. Grace. "Instead, we found that patients from socioeconomically advantaged areas were less likely to use peritoneal dialysis and more likely to use in-centre hemodialysis."

It's unclear why private hospitals in Australia rarely use home dialysis, and why patients from advantaged areas are more likely to use private hospitals.

"More research is required to determine if patients from advantaged areas choose private hospitals knowing they want to dialyse in centre, or whether they attend public hospitals then get directed towards incenter dialysis," added Grace.

Published on : Wed, 25 Jun 2014