

When 2D Imaging Is Insufficient in Understanding Congenital Heart Disease Cases



The result

Holding a <u>real-size replica of a heart</u> and observing the inner heart from all angles enabled cardiologists to view the pathology in a new light. It opened a new dimension in understanding the morphology of the congenital heart defects, and facilitated the team's discussion on procedural planning and management decisions.

Patient 1 (CMVSD): She had previously endured multiple failed attempts at device closure of the complex residual VSDs. Using the 3D replica led to a successful device closure.

Patient 2 (CMVSD): The heart replica provided insightful information about why the previous transcatheter device closure had been unsuccessful. It helped to quickly identify a better treatment option, which was a surgical closure.

Patient 3 (CMVSD): A better understanding of the complexity of the device placement allowed the team to change the treatment option to surgical repair with patch closure and re-endocardialization.

Patient 4 (DORV): The initial plan, based on CT imaging, was to perform VSD closure with baffling to the aortic valve. The 3D cardiac model provided a clearer view of the relationship of the VSD to both great arteries, which led to the decision of a Yasui operation.

Patient 5 (DORV): An aortopulmonary shunt was successfully placed. Future plans involve baffling the VSD from the LV to the aortic valve.

Patient 6 (DORV): A better view on the aortic valve pathway helped defend the decision to proceed with a single ventricle palliation.

What we can conclude

In all six cases, the patient-specific heart replica enhanced the team's understanding of the complex anatomy. In patient 1's case, it was used for a real-time comparison with the angiographic images guiding the intervention. In patients 2 and 3, it provided an in-depth 3D knowledge of the intracavitary anatomy, which greatly reduced the amount of time needed to locate and close the VSDs. The models also helped the team to better grasp the complexities of device closure, the reasons for failure in patient 2, and helped in appropriate patient selection in patient 3.

The additional insight conveyed by the models for preoperative inspection and procedural planning also demonstrates what CT and MRI scans alone fall short in providing. This study shows us that using 3D-printed cardiac models in the treatment of DORV and CMVSD CHDs is invaluable in determining the feasibility of difficult surgical and interventional strategies

Source & Image Credit: Materialise Medical

Published on: Thu, 18 May 2017