

Young Women With Diabetes have Six-Fold Risk of Heart Attack



Women aged 45 years and under with diabetes have a six-fold risk of heart attack, according to research presented at ESC Congress. The study in more than 7 000 women also found that young women who had a heart attack (myocardial infarction, MI) were more likely to be smokers than older women with MI.

"Cardiovascular diseases (CVDs) affect mainly the elderly, but for many years an increase in incidence has been observed in young people as well, regardless of gender," said Professor Hanna Szwed, last author and head of the 2nd Department of Coronary Artery Disease, Institute of Cardiology in Warsaw, Poland. "The World Health Organization estimates that CVDs cause more than 52% of all deaths in women and the number continues to rise. Up to 1% of all heart attacks are in young women."

This study in 7 386 women assessed the effect of risk factors on MI incidence in young women and assessed differences related to age. The epidemiological profile of 1 941 young women (aged ≤45 years) with MI was retrospectively compared with two control groups: 4 275 women aged 63-64 years with MI and 1 170 healthy women (no MI history) aged ≤45 years.

The study used data from three national registries. Data on women with MI came from the Polish Registry of Acute Coronary Syndromes (PL-ACS).² Information on healthy women was from the Multicentre Study of State National Population Health (WOBASZ)³ and National Survey of Risk Factors for Cardiovascular Diseases (NATPOL).⁴

The average age among young MI patients in the study was 42 years (range 21-45 years). Multivariate analysis showed that four out of five classic risk factors were independent predictors of MI in young women. The strongest was diabetes which increased MI risk by six-fold. Arterial hypertension increased risk by four times while hypercholesterolemia tripled risk and current smoking increased risk by 1.6 times. There was no statistical significance for obesity expressed by body mass index (BMI).⁵

Professor Szwed said: "The lack of a correlation with obesity could be because of the overwhelming influence of diabetes in this population. We also found that the risk of MI in young women increased with the number of coexisting factors (Figure 1)."

The researchers found that young healthy women had an average of 1.1 risk factors, while young MI patients had 1.7 and older women with MI had 2.0. Compared to young healthy women, young women with MI more often had arterial hypertension, hypercholesterolemia and diabetes and were current or past smokers (Figure 2).⁶

In the population of older women with MI, the prevalence of nearly all individual risk factors was even greater than in young women with MI.⁷ Smoking was the only risk factor with a greater prevalence in the younger group (48.7% vs. 22.2% $p < 0.0001$ for current smoking, and 65.5% vs. 45.1% $p < 0.0001$ for past smoking) (Figure 2). "We found that the risk factor profile in young women with MI was similar to the older population apart from the greater occurrence of tobacco smoking in young women," said Professor Szwed. "This finding correlates with other research which shows that smoking is a growing problem in young women. This is clearly an area where prevention efforts are needed." Interestingly, young women with MI were more likely than older women to have none of the above-mentioned risk factors (16.5% vs. 8.2% $p < 0.0001$). The most frequent risk factor in this subgroup of patients (both for younger and older women) was a family history of coronary artery disease, with a higher rate in younger compared to older women (11.9% vs. 5.5% $p = 0.0035$).

Professor Szwed concluded: "At present there are not enough global scientific reports focused on the problem of coronary heart disease in young populations, particularly in women. More research is needed into this growing problem to deepen our knowledge, improve prevention efforts and reduce mortality."

Figure 1: Relationship between average number of risk factors and rate of heart attacks in young women according to age





Figure 2: Prevalence of risk factors by age and history of myocardial infarction



NATPOL/WOBASZ: healthy women aged <45 years; PL-ACS: women with history of MI

Source: ESC

Image Credit: Wikimedia Commons

Published on : Mon, 31 Aug 2015